

Role of Bypass Nutrients in Dairy Animals

Anupam Soni and Harsimar Preet Kour*

Livestock Production Management Division, NDRI, Karnal, Haryana, India

Corresponding Author

Harsimar Preet Kour

Email: harsimar0303@gmail.com



OPEN ACCESS

Keywords

Bypass fat, Bypass protein, Dairy animals, protected amino acids, Transition period

How to cite this article:

Soni, A. and Kour, H. P. 2026. Role of Bypass Nutrients in Dairy Animals. *Vigyan Varta* 7 (06): 33-40.

ABSTRACT

Bypass nutrients have emerged as an important nutritional strategy for improving the productivity, health and reproductive efficiency of dairy animals. In high-yielding dairy cows and buffaloes, conventional feeding practices often fail to supply sufficient nutrients because a large proportion of dietary protein and energy is degraded in the rumen before absorption. Bypass nutrients are specially protected feed components that escape ruminal degradation and become available for digestion in the intestine, thereby improving nutrient utilization efficiency. The major bypass nutrients include bypass protein, bypass fat and protected amino acids such as methionine and lysine. Supplementation of bypass protein enhances amino acid availability for milk synthesis, growth and reproductive functions, while bypass fat provides concentrated energy during periods of high metabolic demand, particularly in early lactation. Protected amino acids support milk protein synthesis, liver function and immune response. Feeding bypass nutrients helps reduce negative energy balance, ketosis and other metabolic disorders commonly observed during the transition period. In addition, these nutrients improve milk yield, milk fat percentage, body condition score, conception rate and overall animal performance. Bypass feeding is also beneficial under tropical conditions where poor-quality roughages are widely used. Improved nutrient utilization through bypass supplementation contributes to sustainable dairy farming by reducing nutrient wastage and enhancing feed conversion efficiency. Despite their advantages, factors such as cost, quality control and balanced ration formulation must be considered for effective utilization. Recent advancements in feed processing technologies and rumen-protection techniques are expanding the scope of bypass nutrition in modern dairy

production systems. Overall, bypass nutrients represent an effective approach for maximizing production efficiency, improving animal welfare and increasing profitability in the dairy sector.

INTRODUCTION

Dairy farming plays a vital role in the agricultural economy and nutritional security of many countries, especially India (Patton, 1999; NRC, 2001). High-producing dairy animals require balanced nutrition for maintaining health, reproduction, milk production and overall productivity (Santos *et al.*, 1998). However, conventional feeding practices often fail to meet the nutrient demands of high-yielding dairy cows and buffaloes, particularly during early lactation and transition periods (Grummer, 1995). In ruminants, a significant portion of dietary nutrients is degraded in the rumen before reaching the intestine. This reduces the efficiency of nutrient utilization and may lead to production losses (Chalupa, 1975). To overcome this limitation, the concept of bypass nutrients has gained importance in modern dairy nutrition (NRC, 2001). Bypass nutrients are feed nutrients that escape degradation in the rumen and are digested and absorbed in the intestine (Stern *et al.*, 1994). These nutrients provide a direct supply of high-quality protein, fat and amino acids to the animal, thereby improving productivity and metabolic efficiency (Palmquist & Jenkins, 1980). Feeding bypass nutrients has become an important strategy for enhancing milk yield, milk composition, reproductive performance and health status of dairy animals (Naik *et al.*, 2009).

Concept of Bypass Nutrients

The digestive system of ruminants is unique because feed first enters the rumen, where microbes ferment carbohydrates, proteins and fats (Van Soest, 1994). Although rumen microbes synthesize microbial protein and

vitamins, excessive degradation of nutrients in the rumen may reduce nutrient availability to the animal itself (Stern *et al.*, 1994). Bypass nutrients are specially processed nutrients designed to resist ruminal degradation and become available for digestion in the small intestine (Chalupa, 1975).

The main bypass nutrients used in dairy nutrition include bypass protein, bypass fat and protected amino acids (NRC, 2001). These nutrients help animals meet the high energy and protein requirements during periods of stress, growth and lactation (Grummer, 1995). Bypass feeding is particularly beneficial for high-yielding animals, transition cows, pregnant animals and buffaloes under tropical conditions (Naik *et al.*, 2009).

Importance of Bypass Nutrients in Dairy Animals

Modern dairy animals are genetically superior and capable of producing large quantities of milk (Santos *et al.*, 1998). Such animals require dense nutrient supply beyond what rumen fermentation alone can provide (NRC, 2001). Bypass nutrients improve nutrient efficiency and reduce metabolic stress in dairy animals (Palmquist & Jenkins, 1980). Feeding bypass nutrients helps in increasing milk production, improving milk fat percentage, enhancing fertility and reducing negative energy balance (Grummer, 1995). It also supports immune function and reduces the incidence of metabolic disorders such as ketosis and fatty liver (Drackley, 1999). In tropical countries, where poor-quality roughages are commonly fed, bypass nutrients

help improve the utilization efficiency of available feeds (Naik *et al.*, 2009).

Types of Bypass Nutrients

1. Bypass Protein

Bypass protein, also called undegradable dietary protein (UDP) or rumen undegradable protein (RUP), is the portion of dietary protein that escapes ruminal degradation and reaches the intestine intact (Stern *et al.*, 1994). In the intestine, it is digested into amino acids and absorbed by the animal (NRC, 2001). Common bypass protein sources include heat-treated soybean meal, maize gluten meal, fish meal, cottonseed cake and treated oil cakes (Chalupa, 1975). These protein sources provide essential amino acids required for milk synthesis, tissue repair and reproductive functions (Santos *et al.*, 1998).

2. Bypass Fat

Bypass fat refers to fat sources protected from ruminal degradation (Palmquist & Jenkins, 1980). Normal fats can interfere with rumen microbial activity when fed in excess (Jenkins, 1993). Protected fats are specially processed to avoid rumen fermentation and are digested in the intestine (Palmquist, 1984). Calcium salts of long-chain fatty acids are commonly used as bypass fat supplements (Jenkins & Palmquist, 1984). Bypass fat is an excellent energy source and is highly beneficial during early lactation when animals experience negative energy balance (Grummer, 1995).

3. Protected Amino Acids

Certain amino acids such as methionine and lysine are essential for milk protein synthesis (Schwab, 1996). In normal feeding conditions, these amino acids may be degraded in the rumen before absorption (Stern *et al.*, 1994). Protected amino acids are coated or chemically treated to escape rumen degradation (NRC, 2001). Supplementation of protected amino

acids improves milk yield, milk protein percentage and reproductive efficiency (Schwab *et al.*, 2004).

Mechanism of Action of Bypass Nutrients

Bypass nutrients escape microbial degradation in the rumen and pass directly to the abomasum and small intestine (Chalupa, 1975). There, they are digested enzymatically and absorbed efficiently (Van Soest, 1994). Bypass protein provides essential amino acids directly to body tissues (Santos *et al.*, 1998). Bypass fat increases dietary energy density without disturbing rumen fermentation (Palmquist & Jenkins, 1980). Protected amino acids ensure availability of limiting amino acids required for milk protein synthesis (Schwab, 1996). Overall, bypass nutrients improve feed efficiency and metabolic performance (NRC, 2001).

Role of Bypass Protein in Dairy Animals

Improvement in Milk Production

Milk synthesis requires a continuous supply of amino acids (Santos *et al.*, 1998). Bypass protein provides high-quality amino acids directly to the intestine, increasing milk production and lactation persistency (Stern *et al.*, 1994). High-yielding cows often respond positively to bypass protein supplementation (NRC, 2001). Animals receiving bypass protein generally produce more milk because of improved nitrogen utilization and enhanced mammary gland function (Broderick *et al.*, 2009).

Enhancement of Milk Composition

Bypass protein improves milk protein percentage and solids-not-fat (SNF) content (Santos *et al.*, 1998). Amino acids absorbed from bypass protein are utilized for casein synthesis in milk (Schwab, 1996). Improved milk quality increases the economic value of milk (NRC, 2001). Some studies have also

reported improvement in milk fat percentage due to better energy-protein balance (Naik *et al.*, 2009).

Improvement in Growth and Body Condition

Growing calves and heifers require adequate protein for muscle development and skeletal growth (NRC, 2001). Feeding bypass protein enhances body weight gain and feed conversion efficiency (Stern *et al.*, 1994). In lactating animals, bypass protein helps maintain body condition score by reducing excessive mobilization of body reserves (Grummer, 1995).

Role in Reproduction

Protein deficiency negatively affects reproductive efficiency (Butler, 2000). Bypass protein improves ovarian activity, conception rate and postpartum recovery (Santos *et al.*, 1998). Animals receiving adequate bypass protein show earlier estrus and reduced days open (Butler, 2000). Better reproductive performance results from improved hormone synthesis and energy balance (NRC, 2001).

Role of Bypass Fat in Dairy Animals

Source of Concentrated Energy

Fat contains approximately 2.25 times more energy than carbohydrates (Palmquist & Jenkins, 1980). Bypass fat increases dietary energy density without increasing grain feeding (Grummer, 1995). This is particularly useful during early lactation when feed intake is low (Drackley, 1999). Supplementation of bypass fat helps animals meet energy requirements and reduces negative energy balance (Naik *et al.*, 2009).

Reduction of Ketosis and Metabolic Disorders

High-yielding dairy animals are prone to ketosis during early lactation due to energy

deficiency (Drackley, 1999). Bypass fat provides additional energy and reduces mobilization of body fat reserves (Grummer, 1995). Adequate energy supplementation decreases ketone body formation and helps prevent metabolic disorders such as ketosis and fatty liver syndrome (Duffield, 2000).

Improvement in Milk Yield and Fat Percentage

Bypass fat supplementation often increases milk yield and milk fat content (Palmquist, 1984). Fatty acids absorbed from bypass fat are utilized in milk fat synthesis (Jenkins & Palmquist, 1984). Animals supplemented with bypass fat usually show improved lactation performance and better persistency (Naik *et al.*, 2009).

Improvement in Reproductive Performance

Energy deficiency delays ovarian activity and conception (Butler, 2000). Bypass fat improves energy status and enhances reproductive hormones (Staples *et al.*, 1998). Supplementation is associated with improved conception rate and shorter calving interval (Staples *et al.*, 1998). Certain fatty acids also influence prostaglandin synthesis and embryo survival (Mattos *et al.*, 2000).

Role of Protected Amino Acids

Methionine: Methionine is often the first limiting amino acid in dairy rations (Schwab, 1996). It is essential for milk protein synthesis, liver function and immunity (Osorio *et al.*, 2013). Protected methionine improves milk protein yield and reduces fatty liver risk (Osorio *et al.*, 2013). Methionine also acts as a methyl donor and supports metabolic functions (NRC, 2001).

Lysine: Lysine is another important amino acid for milk production and growth (Schwab *et al.*, 2004). Protected lysine supplementation improves milk yield and protein synthesis

(NRC, 2001). Balanced methionine and lysine nutrition enhances overall production efficiency (Schwab *et al.*, 2004).

Bypass Nutrients During Transition Period

The transition period, extending from three weeks before calving to three weeks after calving, is the most critical phase in dairy animals (Grummer, 1995). Nutrient demands rise sharply while feed intake decreases (Drackley, 1999). Feeding bypass protein and bypass fat during this period helps reduce negative energy balance, improves immunity and lowers the incidence of metabolic diseases (Butler, 2000). Transition cows supplemented with bypass nutrients show better milk production and reproductive performance after calving (Santos *et al.*, 1998).

Bypass Nutrients and Immune Function

Nutritional deficiencies weaken the immune system and increase disease susceptibility (Kehrli *et al.*, 1989). Bypass nutrients support immune function by supplying essential amino acids and energy (NRC, 2001). Adequate protein and energy improve antibody production, leukocyte function and disease resistance (Kehrli *et al.*, 1989). Animals receiving bypass nutrients generally show lower incidence of mastitis, retained placenta and uterine infections (Drackley, 1999).

Bypass Nutrients and Heat Stress Management

Heat stress reduces feed intake and milk production in dairy animals (West, 2003). During hot climates, animals often fail to consume enough nutrients to meet production requirements (West, 2003). Bypass fat provides additional energy without increasing heat production during digestion (Palmquist & Jenkins, 1980). Feeding bypass nutrients during summer helps maintain milk production and body condition under heat stress conditions (Naik *et al.*, 2009).

Sources of Bypass Protein

Soybean meal: Heat-treated soybean meal is widely used as bypass protein and improves milk production (Santos *et al.*, 1998).

Cottonseed cake: Formaldehyde-treated cottonseed cake is commonly used in India as an economical bypass protein source (Walli *et al.*, 1993).

Fish meal: Fish meal contains excellent quality protein and essential amino acids (NRC, 2001). However, its use may be limited by cost and availability.

Maize gluten meal: It is rich in bypass protein and energy and is frequently used in dairy rations (Chalupa, 1975).

Sources of Bypass Fat

Calcium salts of fatty acids are the most common commercial bypass fat supplements used in dairy feeding (Jenkins & Palmquist, 1984).

Prilled fat contains saturated fatty acids processed into granules and is highly digestible for high-yielding dairy animals (Naik *et al.*, 2009).

Hydrogenated vegetable oils are also used as bypass fat supplements (Palmquist, 1984).

Methods of Preparing Bypass Protein

Heat treatment reduces ruminal degradability of proteins and increases bypass value (Chalupa, 1975).

Chemical treatment using formaldehyde protects protein from microbial degradation in the rumen (Walli *et al.*, 1993).

Extrusion technology uses heat and pressure to protect proteins and improve digestibility (NRC, 2001).

Feeding Strategies for Bypass Nutrients

Bypass nutrients should be fed according to production stage, milk yield and body condition (NRC, 2001). High-producing cows require greater supplementation than low-yielding animals (Grummer, 1995). Bypass fat is usually fed at 100–200 g/day in cows, while bypass protein supplementation depends on dietary protein deficiency (Naik *et al.*, 2009). Balanced feeding is essential because excessive supplementation may increase feed cost without additional benefit (Palmquist & Jenkins, 1980).

Economic Benefits of Bypass Nutrients

Supplementation of bypass nutrients increases milk yield and improves reproductive efficiency, leading to higher farm profitability (Naik *et al.*, 2009). Improved feed efficiency reduces wastage of nutrients and lowers production cost per litre of milk (NRC, 2001). Reduced disease incidence and improved fertility also decrease veterinary and breeding expenses (Butler, 2000).

Bypass Nutrients in Buffaloes

Buffaloes are important dairy animals in India and have high milk fat content (Paul & Lal, 2010). Feeding bypass nutrients to buffaloes improves milk production, reproductive performance and body condition (Naik *et al.*, 2009). Studies have shown that bypass fat supplementation in Murrah buffaloes significantly increases milk yield and milk fat percentage during early lactation (Tyagi *et al.*, 2010).

Role in Sustainable Dairy Farming

Efficient nutrient utilization is important for sustainable dairy production (FAO, 2013). Bypass nutrients reduce nitrogen losses in manure and improve feed conversion efficiency (NRC, 2001). Improved nutrient utilization also reduces methane emission

intensity per unit of milk production (Hristov *et al.*, 2013). Thus, bypass feeding contributes to environmentally sustainable dairy farming.

Limitations of Bypass Nutrients

Despite several benefits, bypass nutrients have some limitations. Commercial bypass supplements may be expensive for small farmers (Naik *et al.*, 2009). Improper supplementation can lead to nutrient imbalance and digestive disturbances (NRC, 2001). Quality control and proper ration formulation are essential for obtaining maximum benefit from bypass feeding.

Future Scope of Bypass Nutrition

Advances in feed processing technologies are improving the effectiveness of bypass nutrients (NRC, 2001). Development of rumen-protected vitamins, minerals and amino acids is expanding the scope of bypass nutrition (Schwab *et al.*, 2004). Precision feeding systems and nutrigenomics may further enhance nutrient utilization efficiency in dairy animals. Future research is focused on economical and eco-friendly bypass feed technologies suitable for tropical dairy systems (FAO, 2013).

CONCLUSION

Bypass nutrients play an important role in improving productivity, health and reproductive efficiency of dairy animals. Bypass protein, bypass fat and protected amino acids help overcome the limitations of ruminal nutrient degradation and ensure better nutrient availability to the animal. Feeding bypass nutrients is especially beneficial for high-yielding and transition dairy animals. These nutrients improve milk yield, milk composition, fertility and metabolic health while supporting sustainable dairy farming. Adoption of bypass feeding technologies can significantly enhance dairy profitability and

animal welfare under modern intensive production systems.

REFERENCES

- Broderick, G. A., Stevenson, M. J., & Patton, R. A. (2009). Effect of dietary protein concentration and degradability on lactation performance of dairy cows. *Journal of Dairy Science*, 92(6), 2719–2728.
- Butler, W. R. (2000). Nutritional interactions with reproductive performance in dairy cattle. *Animal Reproduction Science*, 60–61, 449–457.
- Chalupa, W. (1975). Rumen bypass and protection of proteins and amino acids. *Journal of Dairy Science*, 58(8), 1198–1218.
- Drackley, J. K. (1999). Biology of dairy cows during the transition period: The final frontier? *Journal of Dairy Science*, 82(11), 2259–2273.
- Duffield, T. (2000). Subclinical ketosis in lactating dairy cattle. *Veterinary Clinics of North America: Food Animal Practice*, 16(2), 231–253.
- FAO. (2013). Mitigation of greenhouse gas emissions in livestock production: A review of technical options for non-CO₂ emissions. Food and Agriculture Organization of the United Nations, Rome, Italy.
- Grummer, R. R. (1995). Impact of changes in organic nutrient metabolism on feeding the transition dairy cow. *Journal of Animal Science*, 73(9), 2820–2833.
- Hristov, A. N., Oh, J., Firkins, J. L., Dijkstra, J., Kebreab, E., Waghorn, G., et al. (2013). Mitigation of methane and nitrous oxide emissions from animal operations: III. A review of animal management mitigation options. *Journal of Animal Science*, 91(11), 5095–5113.
- Jenkins, T. C. (1993). Lipid metabolism in the rumen. *Journal of Dairy Science*, 76(12), 3851–3863.
- Jenkins, T. C., & Palmquist, D. L. (1984). Effect of fatty acids or calcium soaps on rumen and total nutrient digestibility of dairy rations. *Journal of Dairy Science*, 67(5), 978–986.
- Kehrli, M. E., Nonnecke, B. J., & Roth, J. A. (1989). Alterations in bovine neutrophil function during the periparturient period. *American Journal of Veterinary Research*, 50(2), 207–214.
- Mattos, R., Staples, C. R., & Thatcher, W. W. (2000). Effects of dietary fatty acids on reproduction in ruminants. *Reviews of Reproduction*, 5(1), 38–45.
- Naik, P. K., Saijpal, S., Rani, N., & Dhuria, R. K. (2009). Effect of rumen protected fat on performance of lactating dairy animals. *Indian Journal of Animal Nutrition*, 26(1), 1–12.
- NRC. (2001). Nutrient requirements of dairy cattle (7th rev. ed.). National Academy Press, Washington, DC.
- Osorio, J. S., Ji, P., Drackley, J. K., Luchini, D., & Looor, J. J. (2013). Smartamine M supplementation during the periparturient period alters liver function and inflammation biomarkers in dairy cows. *Journal of Dairy Science*, 96(11), 7120–7133.
- Palmquist, D. L. (1984). Use of fats in diets for lactating dairy cows. In: *Fats in Animal Nutrition* (pp. 357–381). Butterworths, London.

- Palmquist, D. L., & Jenkins, T. C. (1980). Fat in lactation rations: Review. *Journal of Dairy Science*, 63(1), 1–14.
- Patton, S. (1999). *Milk: Its remarkable contribution to human health and well-being*. Transaction Publishers, New Brunswick, NJ.
- Paul, S. S., & Lal, D. (2010). Nutrient requirements and feeding of buffaloes and cattle. *Indian Journal of Animal Sciences*, 80(11), 1065–1074.
- Santos, F. A. P., Huber, J. T., Theurer, C. B., Nussio, L. G., Tarazon, M., & Fish, D. (1998). Response of high producing dairy cows to different protein sources. *Journal of Dairy Science*, 81(1), 192–203.
- Schwab, C. G. (1996). Rumen-protected amino acids for dairy cattle: Progress towards determining lysine and methionine requirements. *Animal Feed Science and Technology*, 59(1–3), 87–101.
- Schwab, C. G., Huhtanen, P., Hunt, C. W., & Hvelplund, T. (2004). Nitrogen requirements of cattle. In: *Protein and Energy Requirements of Ruminants* (pp. 13–70). CAB International.
- Staples, C. R., Burke, J. M., & Thatcher, W. W. (1998). Influence of supplemental fats on reproductive tissues and performance of lactating cows. *Journal of Dairy Science*, 81(3), 856–871.
- Stern, M. D., Bach, A., & Calsamiglia, S. (1994). Alternative techniques for measuring nutrient digestion in ruminants. *Journal of Animal Science*, 72(8), 2256–2276.
- Tyagi, N., Thakur, S. S., & Shelke, S. K. (2010). Effect of bypass fat supplementation on productive and reproductive performance in crossbred cows and buffaloes. *Tropical Animal Health and Production*, 42(8), 1749–1755.
- Van Soest, P. J. (1994). *Nutritional ecology of the ruminant* (2nd ed.). Cornell University Press, Ithaca, NY.
- Walli, T. K., Garg, M. R., & Ravi Kumar, A. (1993). Effect of formaldehyde-treated protein supplements on nutrient utilization in dairy animals. *Indian Journal of Dairy Science*, 46(9), 395–399.
- West, J. W. (2003). Effects of heat stress on production in dairy cattle. *Journal of Dairy Science*, 86(6), 2131–2144.